





Barium-ion tagging for ^{136}Xe double-beta decay studies with EXO



Thomas Brunner for the EXO collaboration
TIPP2014 – June 5, 2014


EXO– a multi-phase approach

The virtues of ^{136}Xe in a large TPC


- **Easy to enrich**: 8.9% natural abundance but can be enriched relatively easily (better than growing crystals)
- **Can be purified** continuously, and reused
- **High $Q_{\beta\beta}$** (2458 keV): higher than most naturally occurring backgrounds
- **Minimal cosmogenic activation**: no long-life radioactive isotopes
- **Energy resolution**: improves using scintillation and charge anti-correlation
- **LXe self shielding**
- Background can be potentially reduced by **Ba^{++} tagging**


Phased approach:

1. EXO-200: 200kg liquid-Xe TPC




2. nEXO: 5-ton liquid Xe TPC with Ba tagging option (SNO lab cryopit)






EXO-200



See poster by Lisa Kaufmann

- 2150 feet depth (1585 mwe)
- Low radioactivity levels:
 - U, Th <100ppb
 - Radon background < 10 Bq/m³
- Liquid Xe TPC (200 kg, 80.6% enriched ¹³⁶Xe)
- Charge and scintillation light readout




Exposure $0\nu\beta\beta$ analysis:
99.8 kg yr

♦ Data
 — Best Fit
 Rn
 - - - LXe bgd
 — n -capture
 - - - ²³²Th (far)
 — Vessel
 — $0\nu\beta\beta$
 — $2\nu\beta\beta$

$T_{1/2}(0\nu) > 1.1 \times 10^{25}$ yr (90%CL) arXiv:1402.6956v1
 $T_{1/2}(2\nu) = 2.165 \pm 0.016$ (stat) ± 0.059 (sys) $\times 10^{21}$ yr
 PRC 89(2014)015502

Enriched Xenon Observatory

- Multi-phase program
- EXO-200, in operation:
 - 200 kg LXe
 - Sensitivity: 100-200 meV
- Multi-ton EXO, R&D underway:
 - 5 ton liquid Xe
 - Sensitivity: 5-30 meV
 - Improved techniques for background suppression and possibly Ba tagging



→ Development of nEXO, a multi-ton scale detector, is well advanced

Barium tagging in EXO

Idea: Perform a background-free measurement by identifying the decay product – Ba-ion tagging

Sensitivity with background $\langle m_\nu \rangle \propto 1 / \sqrt{T_{1/2}^{0\nu\beta\beta}} \propto 1 / (Nt)^{1/4}$


Sensitivity without background $\langle m_\nu \rangle \propto 1 / \sqrt{T_{1/2}^{0\nu\beta\beta}} \propto 1 / (Nt)^{1/2}$

Ba-tagging concept

1. Determine event energy is close to $Q_{\beta\beta}$ (2540keV)
2. Determine position of event
3. Extract decay volume and probe for Barium




Detecting daughter ^{136}Ba provides a “tag” that can discriminate against all background except $2\nu\beta\beta$ decay.






Tagging from Liquid

1. Detect and localize decay (like in EXO200)
2. Send probe in to region of decay
3. Confine the Ba^{++} on probe
4. Remove the probe
5. Identify the barium



Ba⁺ tagging by Resonance Ionization




Concept:

RIS - selective ionization of only one element with lasers



- Move probe close to Ba⁺ ion in LXe
- Attach Ba⁺ ion to probe
- Move probe out of LXe
- Laser-ablate Ba atom from probe
- Laser-ionize Ba⁺ by RIS
- Accelerate Ba⁺ ions and identify by TOF

RIS Ba⁺ tagging at Stanford

Blank regions are when RIS lasers are blocked




RIS and ablated Ba⁺ as well as background ablated ions separated by time-of-flight



nm
1064
553.5
389.7

Ba⁺




Barium tagging in solid xenon (CSU)


Tagging concept

Capture Ba^+ daughter in solid xenon on a probe:


Detect single Ba^+ or Ba on probe by fluorescence:



Solid Xe formed on a cryoprobe in liquid xenon




Barium tagging test apparatus



Successful spectroscopy of Ba-ions in sXe (CSU)


50 ms images of deposit of $\sim 10^5$ Ba⁺ ions in solid xenon
most signal comes in $\sim 5 \mu\text{s}$ before optical pumping occurs

run 18, frame 1 (20130816)




Ba

run 18, frame 2 (20130816)




Ba⁺


run 18, frame 3 (20130816)



Pixels $20\mu\text{m} \times 20\mu\text{m}$



~ 1 in 350 decays from $1P_1$ state are into metastable D states




~ 1 in 4 decays from $2P_{1/2}$ state are into metastable D state

Need repumping lasers to overcome optical pumping and get single Ba/Ba⁺ images with $10^6\times$ or more signal


Barium tagging by Thermal Ionization (IU, TUM)

- Study neutralization of Ba in Xenon Ice.
- Study of desorption of Ba from surfaces.


CARIBU beam at Argonne National Lab
provides radioactive beams of ^{139}Cs or ^{139}Ba



30% transport of Ba ion from Ta surface at 1250 K




Xe Ice on Electrode





General Concept of Ba^{++} Tagging in gas

- Guide Ba^{++} in high pressure Xe inside the TPC (10 bar) to a nozzle
- Extract Ba^{++} with a Xe gas jet into a low pressure chamber
- After nozzle, pump Xe gas away and guide Ba^{++} to identification



Concept of RF-funnel



- Concept of funnel by V. Varentsov
- Conv.-diverging supersonic nozzle
- 301 RF electrodes (0.1 mm thick)
- 0.25 mm electrode spacing
- RF applied to electrodes
- $P_0 = 10 \text{ bar!}$ to 1 mbar in only one stage
- Simulated extraction efficiency of up to 95%
- Xe gas is recaptured by a cryo pump

Pictures of RF-funnel





Photo-etched electrodes with decreasing ID

All components
UHV compatible



2 electrically insulated stacks




Installed RF-funnel during a Xe run



RF-funnel and nozzle

Ion extraction from 10 bar Xe gas



Funnel RF at 2.6 MHz

Current status:

- A Xe or Ar gas jet can be operated at up to 12 bar
- Xe gas can be recovered after an experiment
- Ions can successfully be extracted from high-pressure gas environment

For the future:

- Ion identification
- Determination of extraction efficiency

Conclusion

- Development of nEXO, a multi-ton scale detector, started
- Several groups are working on techniques for Ba-ion extraction from Xe, for the nEXO collaboration
- Successful spectroscopy on Ba-ions in Xe ice (CSU).
- Investigating of Ba-ion properties on surfaces.
- First RIS Ba-ion identification.
- Positive ion extraction from high pressure Xe and Ar gas.

IXe

gXe

The EXO-200 Collaboration



University of Alabama, Tuscaloosa AL, USA - D. Auty, T. Didberidze, M. Hughes, A. Piepke, R. Tsang

University of Bern, Switzerland - S. Delaquis, G. Giroux, R. Gornea, T. Tolba, J-L. Vuilleumier

California Institute of Technology, Pasadena CA, USA - P. Vogel

Carleton University, Ottawa ON, Canada - V. Basque, M. Dunford, K. Graham, C. Hargrove, R. Killick, T. Koffas, F. Leonard, C. Licciardi, M.P. Roza, D. Sinclair

Colorado State University, Fort Collins CO, USA - C. Benitez-Medina, C. Chambers, A. Craycraft, W. Fairbank, Jr., T. Walton

Drexel University, Philadelphia PA, USA - M.J. Dolinski, M.J. Jewell, Y.H. Lin, E. Smith, Y.-R Yen

Duke University, Durham NC, USA - P.S. Barbeau

IHEP Beijing, People's Republic of China - G. Cao, X. Jiang, L. Wen, Y. Zhao

University of Illinois, Urbana-Champaign IL, USA - D. Beck, M. Coon, J. Ling, M. Tarka, J. Walton, L. Yang

Indiana University, Bloomington IN, USA - J. Albert, S. Daugherty, T. Johnson, L.J. Kaufman

University of California, Irvine, Irvine CA, USA - M. Moe

ITEP Moscow, Russia - D. Akimov, I. Alexandrov, V. Belov, A. Burenkov, M. Danilov, A. Dolgolenko, A. Karelin, A. Kovalenko, A. Kuchenkov, V. Stekhanov, O. Zeldovich

Laurentian University, Sudbury ON, Canada - B. Cleveland, A. Der Mesrobian-Kabakian, J. Farine, B. Mong, U. Wichoski

University of Maryland, College Park MD, USA - C. Davis, A. Dobi, C. Hall

University of Massachusetts, Amherst MA, USA - J. Abdollahi, T. Daniels, S. Johnston, K. Kumar, A. Pocar, D. Shy

University of Seoul, South Korea - D.S. Leonard

SLAC National Accelerator Laboratory, Menlo Park CA, USA - M. Breidenbach, R. Conley, A. Dragone, K. Fouts, R. Herbst, S. Herrin, A. Johnson, R. MacLellan, K. Nishimura, A. Odian, C.Y. Prescott, P.C. Rowson, J.J. Russell, K. Skarpaas, M. Swift, A. Waite, M. Wittgen

Stanford University, Stanford CA, USA - J. Bonatt, T. Brunner, J. Chaves, J. Davis, R. DeVoe, D. Fudenberg, G. Gratta, S. Kravitz, D. Moore, I. Ostrovskiy, A. Rivas, A. Schubert, D. Tosi, K. Twelker, M. Weber

Technical University of Munich, Garching, Germany - W. Feldmeier, P. Fierlinger, M. Marino

TRIUMF, Vancouver BC, Canada - J. Dilling, R. Krucken, F. Retière, V. Strickland